

# SURVIVAL OF MALE AND FEMALE *CERASTODERMA GLAUCUM* (BIVALVIA) DURING AERIAL EXPOSURE

K. TARNOWSKA<sup>1</sup>, A. VERNEY<sup>1</sup>, M. WOŁOWICZ<sup>2</sup>, J.P. FERAL<sup>1</sup>, A. CHENUIL<sup>1\*</sup>

<sup>1</sup>IMBE (Mediterranean Institute of Biodiversity and marine and continental Ecology) - UMR 7263, CNRS  
Université Aix-Marseille, France

<sup>2</sup>Laboratory of Estuarine Ecology, Institute of Oceanography, University of Gdańsk, Al. Piłsudskiego 46, 81-378 Gdynia, Poland  
\* Corresponding author: anne.chenuil@imbe.fr

AERIAL EXPOSURE  
*CERASTODERMA GLAUCUM*  
SEX-BIASED DISPERSAL

**ABSTRACT.** – The lagoon cockle, *Cerastoderma glaucum* (Poiret, 1789) usually inhabits small, often isolated, water basins. Dispersal among these basins can take place overland (human mediated) or by migratory birds, when cockles are attached to their bodies. It requires the capacity of the cockle to survive aerial exposure. The time of survival of cockles in air was tested. The LT<sub>50</sub> was of 80 h at 15 °C and 43 h at 22 °C. There was no significant difference in survival time neither among size classes, nor between males and females. The similarity between male and female mortality curves and the strong differences between the results coming from two thermal conditions suggest that the experiment was reliable.

## INTRODUCTION

The lagoon cockle *Cerastoderma glaucum* (Lamellibranchia: Cardiidae) is a euryhaline and eurythermic (Rygg 1970) bivalve present across Europe from the Caspian to the Baltic Seas (Brock 1979). The distribution area of *C. glaucum* even if wide, is very fragmented. It usually inhabits isolated or semi-isolated, shallow, non-tidal biotopes, e.g., brackish lagoons, estuaries, bays and lakes. It was previously thought that the main limiting abiotic factor for this cockle was its intolerance to air exposure provoked by tides (Russell 1971, 1972). However, apparently, the nature of sediment is more important than the direct influence of tides. This species never occurs on bottoms with loose sediment structure typical of regions exposed to tides, currents and waves (Brock 1979).

The pelagic larval stage of the lagoon cockle is restricted to 1-2 weeks (Lauckner 1972, Kingston 1974, Wołowicz 1987). As the distribution areas of this species are often separated by unsuitable habitats (insularity), alternative dispersal modes necessarily allowed the founding of such populations. *C. glaucum* is probably not dispersed by man for aquaculture purposes, because it only has a marginal commercial value (Arjonilla *et al.* 1994). However, it may be transported as a result of anthropogenic activities, such as constructing canals connecting separated basins, or pumping water from one water basin and releasing it into another (e.g. for the needs of power plants) [for a freshwater bivalve, *Dreissena polymorpha* (Pallas, 1771) Quaglia *et al.* 2008]. Furthermore, the spat of *C. glaucum* sticks to the surface of submerged plants, and juveniles climb and attach to substrates with their byssal gland (Pearson 2003, Reise 2003). They may therefore be transported among water basins with these plants attached to the engines of small trailerable boats or to other recreational equipment (Quaglia *et al.* 2008). More-

over, migrating birds seem a plausible vector provoking gene flow (discussed in Tarnowska *et al.* 2010). The role of birds in the dispersal of *C. glaucum* has been previously evoked by many authors (Boyden & Russell 1972, Rose 1972, Gasse *et al.* 1987, Spencer & Patchett 1997, Reise 2003, Nikula 2008). Interestingly, Cadée (1995) suggested a possibility of internal transport of adult cockle, as a specimen of *Cerastoderma edule* (Linnaeus, 1758) was observed alive after regurgitation by a herring gull and found in the act of burrowing again in the tidal flat. However, for mollusks external transport seems to be more common (Wesselingh *et al.* 1999). Adult bivalves may be dispersed attached to birds feet (e.g., Darwin 1878, Kew 1893, Boycott 1936, Rees 1965, Green & Figuerola 2005). Larvae and spat may also be transported externally, for example on birds plumage (Green & Figuerola 2005 and references therein). Additionally, spat and juveniles of *C. glaucum* may be transported by birds together with plants (Reise 2003). Many bird species perform regular migrations between the southern and the northern parts of Europe. They rest and feed in lagoons and estuaries which are often inhabited by *C. glaucum* (Wołowicz 1991). Any overland transport or the transport of cockles attached to birds' feet would require resistance to aerial exposure.

Recent genetic studies of *C. glaucum* populations from almost the whole distribution area revealed very different genetic structure patterns between mitochondrial DNA and microsatellite markers (Tarnowska *et al.* 2010). In mitochondrial DNA, contrary to the nuclear DNA, some geographical discontinuities were found in genetic structure. As mitochondrial DNA is transmitted in the maternal lineage and nuclear DNA is transmitted by both parents (Chenuil 2006), such differences suggest migration rate differences between males and females. It seems unlikely that a male and a female larva behave or even survive differently. However, adult males and females often display

different physiological conditions (thus different survival rates facing stress) due to differences in reproductive efforts in animals (Tarnowska *et al.* 2009). Gender-biased dispersal could hypothetically be driven by gender differences in survival during aerial exposure and starvation involved with passive dispersal mechanisms (*e.g.*, transport by migrating birds or human activity). Gender differences in survival rates, susceptibility to diseases and life span are not rare phenomena (*e.g.*, in human: Olivetti *et al.* 1995, Macintyre *et al.* 1996, Arbuckle 2006) and are even theoretically expected, for instance, in hybrid zones according to sex determinism mechanisms (*e.g.*, Haldane 1922, Chenuil *et al.* 2004). A recent genetic study on the lagoon cockle revealed a striking similarity in the mitochondrial marker of cockles from the Iberian Peninsula and the Baltic Sea (Tarnowska *et al.* 2010). Some waders (Charadriiformes) that undergo particularly long migratory flights use the East Atlantic Flyway, which is a major route for long-distance transport between these two regions (Sánchez *et al.* 2006, Sánchez *et al.* 2007). This suggested that long distance dispersal *via* migrating birds and gender differences in survival during this transport may together explain the particular genetic structure observed in the lagoon cockle at mitochondrial and nuclear markers. Theoretically, even if long distance dispersal occurs for a single individual, it may enable its genotype (or mitochondrial haplotype) to invade an already settled population due to genetic drift or natural selection. This hypothesis led to this experimental study.

## MATERIALS AND METHODS

Cockles used for the experiment were sampled on 8<sup>th</sup> June 2010 in the Berre Lagoon (Plage du Jaï) on the French coast of the Mediterranean Sea (43°24'N, 5°08'E). After sampling, the 84 cockles were transported to the laboratory (30 minutes car trip) in sea water at the same temperature as the sampling site (22 °C). The specimens were sampled from the surface of the sediment. The cockles were divided into two lots, in a way respecting similar size distributions. They were placed on a dry tray, on a laboratory table, in the same orientation, the distance among specimens being of a few centimeters. The first lot was placed at 22 ± 1 °C (exposed to indirect sunlight during the day, in a laboratory room with air conditioning) and the second at 15 ± 1 °C (in the dark). The light conditions differed because of the laboratory conditions constraints. The air temperature was controlled. The number of dead individuals was controlled at more or less regular time intervals. Cockles were considered dead when the shell was slightly open and when applying a contact to the shell did not provoke its closing due to the action of adductor muscles. Each time a cockle was found dead the duration of the survival in air was recorded. The LT<sub>50</sub>, which is the exposure time required to kill 50 % of the test population, was calculated for each temperature. Then shell length, width and height were measured with a slide caliper accurate to 0.01 mm.

The dead cockles were placed individually in small plastic bags filled with sea water at 4 °C and stored until observation (maximum 2 days). Gonad tissue (equivalent to a volume of a few microliters) was spread on a glass slide (those classically used for histology) and the slide was placed on a heating plate at 50 °C for a few seconds. A few drops of toluidine blue were spread at the slide for coloration and the excess of it was rinsed for a few seconds under distilled water flow. The sex of each individual was then determined under an optical microscope according to a reliable protocol, since eggs and sperm are distinguishable regardless of gonad development stage (Wołowicz 1987, 1991).

The exact test of Fisher was applied to test the significance of differences in the sex ratio. The 95 % confidence intervals were calculated for the proportion of females (Newcombe 1998). Other statistical analyses were performed using XLstat v.2011. The normality of the data distribution was checked using the Shapiro-Wilk test (significance level 5 %). The significance of the correlations of the survival time in air and every dimension of the shell (length, width and height) was checked with the Spearman (non-parametric) test. The effects of sex and size (three size groups chosen *a priori* attributing equal number of individuals to each group) on survival time in air were investigated with the Kruskal-Wallis non-parametric test.

## RESULTS

The proportion of females was 0.37 (± 0.1), but the females to males ratio did not significantly differ from 1: 1 ( $p > 0.05$ ). The observed survival time data did not have a normal distribution. The temperature of the room strongly influenced survival time, which was significantly longer at 15 °C than at 22 °C ( $p < 0.0001$ ) (Fig. 1). The LT<sub>50</sub> was 80 h at 15 °C and 43 h at 22 °C. The mortality exceeded 80 % after about 100 h at 15 °C and after about 50 h at 22 °C. No significant difference was found between the survival time of males and females in both samples ( $p = 0.644$  at 15 °C,  $p = 0.540$  at 22 °C) and the whole curves were very similar (Fig. 1). The survival time at both temperatures slightly diminished when cockle size increased, but neither the correlations ( $p = 0.346$  and 0.635 respectively at 15 °C and 22 °C) nor the Kruskal-Wallis tests ( $p = 0.838$  and 0.628 respectively at 15 °C and 22 °C) were significant.

## DISCUSSION

### *Survival time and the role of air temperature*

In our study the LT<sub>50</sub> of *C. glaucum* during the aerial exposure was of 80 h (more than 3 days) at 15 °C and




Fig. 1. – Mortality of cockles during the aerial exposure for each sex at 15 °C and 22 °C.

of 43 h (less than 2 days) at 22 °C. These durations were slightly shorter than the ones found by Boyden (1972):  $LT_{50}$  of 87 h (almost 4 days) at 15 °C and of 63 h (almost 3 days) at 20 °C. Comparisons among these studies are difficult to perform in a reliable way since numerous conditions were different. In particular, our study did not control and did not record humidity values, and light conditions were not identical between the two temperature conditions. Survival time of the common cockle, *Cerastoderma edule* was longer than that of *C. glaucum*, reaching 129 h at 15 °C and 89 h at 20 °C (Boyden 1972). This is because some mollusks, especially those inhabiting intertidal regions, like *C. edule*, have a capacity to air-breathe (Boyden 1972 and references therein). *C. glaucum* cannot air-breathe (Boyden 1972). Dance (1958) reported that a freshwater mussel survived 12 months out of water.

The bivalves which cannot air-breathe initially utilize the oxygen dissolved in the water present in the shell. Afterwards, the conditions become anoxic. The ability to survive in anoxic conditions or low oxygen contents varies among bivalves reaching 2 weeks for *Nucula sulcata* (Bronn, 1831) (Taylor *et al.* 1995), 5-17 days for *Nuculoma tenuis* (Montagu, 1808) (in water with extremely low oxygen content) (Moore 1931) or 5 days for *Modiolus demissus* (Dillwyn) (in nitrogen) (Lent 1968).

The principal physiological means of resisting to both aerial exposure and hypoxia is metabolic rate decrease thus energy saving (Storey & Storey 1990). Prolonged aerial exposure provokes tissue hypoxia and metabolic processes leave the aerobic pathways for the anaerobic ones (Ahmad & Chaplin 1984, Eertman & De Zwaan 1994). In bivalves, this leads to end products (*e.g.*, short-chained organic acids or alcohols) that can be reused or easily removed from the cells (Oeschger 1990, De Zwaan & Eertman 1996).

Another factor influencing survival in air is also tolerance to desiccation. *C. glaucum* may survive the loss of 33 % of water from the tissues. Air temperature is crucial during aerial exposure. In the lagoons inhabited by *C. glaucum*, like the Berre Lagoon, where the cockles were sampled, the temperature may exceed 30 °C in the summer (Stora *et al.* 1995, Gouze *et al.* 2005). This species is adapted to these extreme conditions. However, during aerial exposure, temperature increase causes an increase in evaporation of water from the body. Cockles are closed when exposed to air, which limits evaporation (Boyden 1972). But high temperatures may also increase metabolic activity by increasing oxygen demand (Clarke 1991, in *C. glaucum*: Tarnowska *et al.* 2009).

A longer survival of bivalves subject to aerial exposure or hypoxia at lower temperature was reported previously (Matthews & McMahon 1999). In a freshwater bivalve, *Corbicula fluminea* (O.F. Müller, 1774) subject to periodic emersion in its natural habitat, the ranges of median tolerance to aerial exposure were: 23.8-24.9 h at 35 °C, 71.4-78.2 h at 25 °C, and 248.5-341.6 h at 15 °C (Byrne *et al.* 1988). Apart from high temperatures, environments rich in hydrogen sulphide, and high or low salinity can all negatively affect air survival (Shumway *et al.* 1983, Veldhuizen-Tsoerkan *et al.* 1991). Anoxic and aerial survival were demonstrated to provide early warning indicator of contaminant induced stress (De Zwaan & Eertman 1996).

#### *The influence of size on aerial exposure survival time*

We did not evidence a significant effect of size on survival during aerial exposure. Both our study and the previous one (Boyden 1972) were performed on adult individuals. In *Corbicula fluminea* (Bivalvia) larger individuals were proved to lose water at a slower rate and survived longer in air (at 25 °C and 35 °C) than smaller ones

(Byrne *et al.* 1988). Larger individuals were more tolerant to anoxia than smaller ones in other bivalves [Borsa *et al.* 1992 for *Ruditapes decussatus* (Linnaeus, 1758), Matthews & McMahon 1999 for *Dreissena polymorpha*]. However, the pattern may be inverted with larger specimens being less tolerant of anoxia than smaller ones (Matthews & McMahon 1999 for *C. fluminea*). One of the main sources of anaerobic metabolic substrate in mollusks is glycogen (De Zwaan 1983). Therefore, lower tolerance of larger specimens to anoxia might be due to the reduction of tissue glycogen level correlated with reproductive cycle (Tarnowska *et al.* 2009 and references therein).

In general, smaller, younger bivalves appeared more sensitive to other types of environmental stress, such as intoxication (*e.g.*, Ringwood 1993, Markich 2003). However, in *Mytilus edulis* (Linnaeus, 1758) the resistance to intoxication with copper declined gradually, being significantly correlated with shell length (Hoare & Davenport 1994).

#### ***The influence of sex on the aerial exposure survival time***

No difference was recorded in the survival time in air between males and females. Although the absence of significant difference does not prove that survival is the same among sexes, curves are rather smooth and similar, which can be compared to the conspicuous differences observed among temperature conditions in the same Figure (Fig. 1), and suggests that the sample sizes provided a reasonable statistical power. In previous studies, the females of bivalves were often suggested to be more fragile to various perturbations than males. Stress caused by invasive species *Dreissena* provoked an increased mortality in females in comparison to males in the native bivalve *Lampsilis radiata* (Gmelin, 1791). It was considered to be due to the fact that in female stresses related to breeding added to stresses incurred from *Dreissena* encrustation produced lethal effects (Haag *et al.* 1991). The resistance of females to aerial exposure could decrease during gonad maturation and spawning, as it demands a lot of energy. The gonad development stages were not taken into account in this study. We did not trust gonadic stages inferred from our data, because the stress provoked by aerial exposure may provoke immediate spawning. Reproduction is less seasonal in the Berre lagoon than in other investigated areas (Tarnowska *et al.* 2009) thus this location may not be the most susceptible to reveal sex differences due to reproductive effort.

#### ***The consequences of the results for gene flow***

The survival times of *C. glaucum* in air, although shorter than found by Boyden (1972), are long enough to enable a long-distance dispersal of cockles *via* birds. Theoretically, the dispersal distances may be quite impor-

tant, as for example a duck flies at 60-78 km h<sup>-1</sup> (Welham 1994). Moreover, most bird migrations are at an altitude ranging from 150 m to 600 m, although they may also exceed several thousand meters (Williams 1950). The air temperature is known to decrease with altitude, which is favorable for bivalves survival during birds mediated transport. On the other hand, in our experiment the cockles were placed in order to avoid airflow. The cockles attached to migrating birds are exposed to a high airflow rate likely to accelerate desiccation and death.

Although we confirmed that the time of survival in air is long enough for such a transport, the present study does not support differences in survival times between males and females, which may have explained the different patterns of genetic structure between mitochondrial and nuclear DNA revealed by previous studies (Tarnowska *et al.* 2010). We argued above that absence of statistical significance appeared unlikely to be caused by insufficient sample sizes. However, the survival time in air was studied only on adult individuals. Eggs and juveniles can be transported by birds as well. Their physiology is different than the one of adults and the existence of differences between the two sexes in the aerial exposure survival at these stages cannot be excluded. We collected cockles from the surface of the sediment (not the buried ones), which are more prone to infection by trematodes (Bowers *et al.* 1996). However, birds' feet or necks do not burry into the sediment, so surface sampling is not biased relative to our assumption (bird dispersal). A differential survival among genders during dispersal is not the only possible cause of dispersal differences. In some bivalves, behavior differs between sexes. In *Macoma balthica* (Linnaeus, 1758) females prevailed among the crawlers, compared to the buried individuals (Mouritsen 1997). It is also possible that mitochondrial DNA does not reflect simply neutral processes, but that this marker is subject to natural selection. If mitochondrial haplotypes are differentially adapted to given environmental conditions, then the geographical (or spatial) distribution of mitochondrial genetic diversity does not reflect gene flow (Ballard & Whitlock 2004).

To conclude, in the present study we revealed the influence of air temperature on the survival time of cockles. However, we could not prove that the differences in genetic structures revealed by nuclear and mitochondrial markers were provoked by survival differences to aerial exposure (*i.e.*, birds-mediated transport). The hypothesis of differences among sexes in aerial survival cannot be ruled out. Similar tests could be carried out in other locations, where reproduction is more seasonal, and using more individuals of different sizes. We indicated the non-significant p-values of our tests to allow future meta-analyses to be carried out by computing Fisher's combined probability test. Alternative hypotheses to explain the genetic observations are that (i) differences between the effective dispersal of male and female are provoked by



other mechanisms than aerial survival time of adults (*e.g.*, different behavior) or (ii) the differences between the results revealed by the two genomes reveal an effect of natural selection on mitochondrial DNA or nuclear DNA.

ACKNOWLEDGEMENTS. – We would like to thank C Marschal for his help during microscopic analysis. We acknowledge an anonymous reviewer for corrections to English.

## REFERENCES

- Ahmad TA, Chaplin AE 1984. Anaerobic metabolism of bivalve molluscs during exposure to air. *Biochem Syst Ecol* 12(1): 85-88.
- Arbuckle TE 2006. Are there sex and gender differences in acute exposure to chemicals in the same setting? *Environ Res* 101: 195-204.
- Arjonilla M, Forja JM, Gomez-Parra A 1994. Sediment analysis does not provide a good measure of heavy metal bioavailability to *Cerastoderma glaucum* (Mollusca: Bivalvia) in confined coastal ecosystems. *Bull Environ Contam Toxicol* 52: 810-817.
- Ballard JWO, Whitlock MC 2004. The incomplete natural history of mitochondria. *Mol Ecol* 13: 729-744.
- Borsa P, Jouselin Y, Delay B 1992. Relationships between allozymic heterozygosity, body size, and survival to natural anoxic stress in the palourde *Ruditapes decussatus* L. (Bivalvia: Veneridae). *J Exp Mar Biol Ecol* 155: 169-181.
- Bowers EA, Bartoli P, Russell-Pinto F, James BL 1996. The metacercariae of sibling species of *Meiogymnophallus*, including *M. rebecqui* comb. nov. (Digenea: Gymnophallidae), and their effects on closely related *Cerastoderma* host species (Mollusca: Bivalvia). *Parasitol Res* 82: 505-510.
- Boycott AE 1936. The habitats of fresh-water Mollusca in Britain. *J Anim Ecol* 5: 116-186.
- Boyden CR 1972. The behaviour, survival and respiration of the cockles *Cerastoderma edule* and *C. glaucum* in air. *J Mar Biol Ass UK* 52: 661-680.
- Boyden CR, Russell PJC 1972. Distribution and habitat range of brackish water cockle (*Cardium (Cerastoderma) glaucum*) in British Isles. *J Anim Ecol* 41: 719-734.
- Brock V 1979. Habitat selection of two congeneric bivalves, *Cardium edule* and *C. glaucum* in sympatric and allopatric populations. *Mar Biol* 54: 149-156.
- Byrne RA, McMahon RF, Dietz TH 1988. Temperature and relative humidity effects on aerial exposure tolerance in the freshwater bivalve *Corlicula fluminea*. *Biol Bull* 175: 253-260.
- Cadée GC 1995. Birds as producers of shell fragments in the Wadden Sea, in particular the role of the Herring Gull. *Géobios MS* 18: 77-85.
- Chenuil A 2006. Choosing the right molecular genetic markers for studying biodiversity: from molecular evolution to practical aspects (review). *Genetica* 127: 101-120.
- Chenuil A, Crespin L, Pouyaud L, Berrebi P 2004. Autosomal differentiation between males and females in hybrid zones: a first report from *Barbus barbus* and *Barbus meridionalis* (Cyprinidae). *Heredity* 93: 128-134.
- Clarke A 1991. What is cold adaptation and how should we measure it? *Am Zool* 31(1): 81-92.
- Dance SP 1958. Drought resistance in an African freshwater bivalve. *J Conchol* 24: 281-283.
- Darwin C 1878. Transplantation of shells. *Nature* 18: 120-121.
- De Zwaan A 1983. Carbohydrate catabolism in bivalves. In Hochachka PW ed, The Mollusca, Vol 1, metabolic biochemistry and molecular biomechanics. Academic Press, Orlando.
- De Zwaan AB, Eertman RHM 1996. Anoxic or aerial survival of bivalves and other euryoxic invertebrates as a useful response to environmental stress- a comprehensive review. *Comp Biochem Physiol C* 113: 299-312.
- Eertman RHM, De Zwaan AB 1994. Survival of the fittest: resistance of mussels to aerial exposure. In Kramer KJM ed, Biomonitoring of coastal waters and estuaries. CRC Press, Boca Raton.
- Gasse F, Fontes JC, Plaziat JC, Carbonel P, Kaczmarek I, de Deckker P, Soulié-Marsche I, Callot Y, Dupeuble PA 1987. Biological remains, geochemistry and stable isotopes for the reconstruction of environmental and hydrological changes in the Holocene lakes from North Sahara. *Palaeogeogr Palaeoclimatol Palaeoecol* 60: 1-46.
- Gouze E, Raimbault P, Garcia N 2005. Ecological survey of the Berre Lagoon. Chemical analysis. Report, GIPREB, COM, Univ Méditerranée, France, 21 p.
- Green A, Figuerola J, 2005. Recent advances in the study of long-distance dispersal of aquatic invertebrates via birds. *Div Distr* 11: 149-156.
- Haag WR, Berg DJ, Garton DW 1991. *Dreissena polymorpha* colonies encrusting native unionid bivalves produce species-specific and sex-specific effects. *J Shellfish Res* 10: 259.
- Haldane JBS 1922. Sex ratio and unisexual sterility in hybrid animals. *J Genet* 12: 101-109.
- Hoare K, Davenport J 1994. Size-related variation in the sensitivity of the mussel, *Mytilus edulis*, to copper. *J Mar Biol Ass UK* 74: 971-973.
- Kew HW 1893. The dispersal of shells. An inquiry into means of dispersal possessed by fresh-water and land Mollusca. London, Kegan Paul, Trench, Trubner and Do., Ltd.
- Kingston P 1974. Some observations on the effects of temperature on the growth of *Cardium edule* and *Cardium glaucum* larvae in the laboratory. *J Mar Biol Ass UK* 54: 309-317.
- Lauckner G 1972. On the taxonomy, ecology and physiology of *Cardium edule* L. and *Cardium lamarcki* Reeve. PhD thesis, Univ Kiel, Germany, 261 p.
- Lent CM 1968. Air gaping by the ribbed mussel, *Modiolus demissus* (Dillwyn). Effects and adaptive significance. *Biol Bull* 134: 60-73.
- Macintyre S, Hunt K, Sweeting H 1996. Gender differences in health: are things really as simple as they seem? *Soc Sci Med* 42: 617-624.
- Markich SJ 2003. Influence of body size and gender on valve movement responses of a freshwater bivalve to uranium. *Environ Toxicol* 18: 126-136.
- Matthews MA, McMahon RF 1999. Effects of temperature and temperature acclimation on survival of zebra mussels (*Dreissena polymorpha*) and Asian clams (*Corbicula fluminea*) under extreme hypoxia. *J Molluscan Stud* 65: 317-325.
- Moore HB 1931. The muds of the Clyde sea area. III. Chemical and physical conditions; rate and nature of sedimentation; and fauna. *J Mar Biol Ass UK* 17: 325-358.
- Mouritsen KN 1997. Crawling behaviour in the bivalve *Macoma balthica*: the parasite- manipulation hypothesis revisited. *Oikos* 79: 513-520.
- Newcombe RG 1998. Two-sided confidence intervals for the single proportion: comparison of seven methods. *Stat Med* 17: 857-872.

- Nikula R 2008. Phylogeography and hybrid swarms: history of brackish water bivalve diversity in North European marginal seas. PhD thesis, Univ Helsinki, Finland, 35 p.
- Oeschger R 1990. Long-term anaerobiosis in sublittoral marine invertebrates from the western Baltic sea: *Halicryptus spinulosus* (Priapulida), *Astarte borealis* and *Arctica islandica* (Bivalvia). *Mar Ecol-Prog Ser* 59: 133-143.
- Olivetti G, Giordano G, Corradi D, Melissari M, Lagrasta C, Gambert SR, Anversa P 1995. Gender differences and aging: effects on the human heart. *J Am Coll Cardiol* 26: 1068-1079.
- Pearson C 2003. The genetic population structure of the lagoon specialists *Nematostella vectensis*, *Cerastoderma glaucum* and *Gammarus insensibilis* from populations along the southern and eastern coasts of the United Kingdom. PhD thesis, Univ Southampton, United Kingdom, 229 p.
- Quaglia F, Lattuada L, Mantecca P, Bacchetta R 2008. Zebra mussels in Italy: where do they come from? *Biol Invasions* 10: 555-560.
- Rees WJ 1965. The aerial dispersal of mollusca. *Proc Malac Soc Lond* 36: 269-282.
- Reise K 2003. Metapopulation structure in the lagoon cockle *Cerastoderma lamarcki* in the northern Wadden Sea. *Helgol Mar Res* 56: 252-258.
- Ringwood AH 1993. Age-specific differences in cadmium sensitivity and bioaccumulation in bivalve mollusks. *Mar Environ Res* 35: 35-39.
- Rose KD 1972. A mollusc new to Lake Birket Qarun, Egypt. *Nautilus* 85: 141-143.
- Russell PJC 1971. A reappraisal of the geographical distributions of the cockles *Cardium edule* L. and *C. glaucum* Bruguière. *J Conchol* 27: 225-234.
- Russell PJC 1972. Biological studies on *Cardium glaucum*, based on some Baltic and Mediterranean populations. *Mar Biol* 16: 290-296.
- Rygg B 1970. Studies on *Cerastoderma edule* (L.) and *Cerastoderma glaucum* (Poiret). *Sarsia* 43: 65-80.
- Sánchez MI, Green AJ, Amat F, Castellanos EM 2007. Transport of brine shrimps via the digestive system of migratory waders: dispersal probabilities depend on diet and season. *Mar Biol* 151: 1407-1415.
- Sánchez MI, Green AJ, Castellanos EM 2006. Internal transport of seeds by migratory waders in the Odiel marshes, southwest Spain: consequences for long-distance dispersal. *J Avian Biol* 37: 201-206.
- Shumway SE, Scott TM, Shick MJ 1983. The effects of anoxia and hydrogen sulphide on survival, activity and metabolic rate in the coot clam, *Mulina lateralis* (Say). *J Exp Mar Biol Ecol* 71: 135-146.
- Spencer JE, Patchett PJ 1997. Sr isotope evidence for a lacustrine origin for the upper Miocene to Pliocene Bouse Formation, lower Colorado River trough, and implications for timing of Colorado Plateau uplift. *Geol Soc Am Bull* 109: 767-778.
- Stora G, Arnoux A, Galas M 1995. Time and spatial dynamics of Mediterranean lagoon macrobenthos during an exceptionally prolonged interruption of freshwater inputs. *Hydrobiologia* 300-301(1): 123-132.
- Storey KB, Storey JM 1990. Metabolic rate depression and biochemical adaptation in anaerobiosis, hibernation and estivation. *Quart Rev Biol* 65: 145-174.
- Tarnowska K, Chenuil A, Nikula R, Féral JP, Wołowicz M 2010. Complex genetic population structure of bivalve (*Cerastoderma glaucum*) living in highly fragmented lagoon habitat. *Mar Ecol-Prog Ser* 406: 173-184.
- Tarnowska K, Wołowicz M, Chenuil A, Féral JP 2009. Comparative studies on the morphometry and physiology of European populations of the lagoon specialist *Cerastoderma glaucum* (Bivalvia). *Oceanologia* 51(3): 1-22.
- Taylor AC, Davenport J, Allen JA 1995. Anoxic survival, oxygen consumption and haemocyanin characteristics in the protobranch bivalve *Nucula sulcata* Bronn. *Comp Biochem Physiol A* 112A(2): 333-338.
- Veldhuizen-Tsoerkan MB, Holwerda DA, Zandee DI 1991. Anoxic survival time and metabolic parameters as stress indices in sea mussels exposed to cadmium or polychlorinated biphenyls. *Arch Environ Contam Toxicol* 20: 259-265.
- Welham CVJ 1994. Flight speeds of migrating birds: a test of maximum range speed predictions from three aerodynamic equations. *Behav Ecol* 5: 1-8.
- Wesselingh FP, Cadée GC, Renema W 1999. Flying high: on the airborne dispersal of aquatic organisms as illustrated by the distribution histories of the gastropod genera *Tryonia* and *Planorbarius*. *Geologie En Mijnbouw* 78: 165-174.
- Williams GG 1950. Weather and spring migration. *Auk* 67: 52-65.
- Wołowicz M 1987. Larval development of *Cardium glaucum* and *C. hauniense* (Bivalvia) from the Gdansk Bay. *Pol Arch Hydrobiol* 34: 107-117.
- Wołowicz M 1991. Geographical differentiation of *Cerastoderma glaucum* Bruguière (Bivalvia) populations. Hypotheses concerning the origin and migration pathways. Univ Gdańsk, Gdańsk, Poland, 151 p.

Received October 24, 2011  
 Accepted January 16, 2012  
 Associate Editor: O Verneau